

Grade 7		
PA Common Core Standard	PA Eligible Content	Common Core State Standard
2.1.7.D.1 Analyze proportional relationships and use them to model and solve real-world and mathematical problems.	M07.A-R.1.1.1 Compute unit rates associated with ratios of fractions, including ratios of lengths, areas and other quantities measured in like or different units. Example: If a person walks 1/2 mile in each 1/4 hour, compute the unit rate as the complex fraction 1/2 / 1/4 miles per hour, equivalently 2 miles per hour.	<b>7.RP.1</b> Compute unit rates associated with ratios of fractions, including ratios of lengths, areas and other quantities measured in like or different units. For example, if a person walks 1/2 mile in each 1/4 hour, compute the unit rate as the complex fraction 1/2/1/4 miles per hour, equivalently 2 miles per hour.
	_	<ul> <li>7.RP.2. Recognize and represent proportional relationships between quantities.</li> <li>a. Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin.</li> </ul>
	M07.A-R.1.1.3 Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships.	<ul> <li>7.RP.2. Recognize and represent proportional relationships between quantities.</li> <li>b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships.</li> </ul>
	<b>M07.A-R.1.1.4</b> Represent proportional relationships by equations. <i>Example: If total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as <math>t = pn</math>.</i>	<ul> <li>7.RP.2. Recognize and represent proportional relationships between quantities.</li> <li>c. Represent proportional relationships by equations. For example, if total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn.</li> </ul>
	<b>M07.A-R.1.1.5</b> Explain what a point (x, y) on the graph of a proportional relationship means in terms of the situation, with special attention to the points (0, 0) and (1, r) where r is the unit rate.	<b>7.RP.2.</b> Recognize and represent proportional relationships between quantities. <b>d.</b> Explain what a point (x, y) on the graph of a proportional relationship means in terms of the situation, with special attention to the points (0, 0) and (1, r) where r is the unit rate.

Updated 5.18.2012 Page 1 of 9



Grade 7		
PA Common Core Standard	PA Eligible Content	Common Core State Standard
	M07.A-R.1.1.6 Use proportional relationships to solve multi-step ratio and percent problems.  Examples: simple interest, tax, markups and markdowns, gratuities and commissions, fees, percent increase and decrease.	<b>7.RP.3.</b> Use proportional relationships to solve multistep ratio and percent problems. <i>Examples: simple interest, tax, markups and markdowns, gratuities and commissions, fees, percent increase and decrease, percent error.</i>
<b>2.1.7.E.1</b> Apply and extend previous understandings of operations with fractions to operations with rational numbers.	and subtract rational numbers, including real-world	<ul> <li>7.NS.1. Apply and extend previous understandings of addition and subtraction to add and subtract rational numbers; represent addition and subtraction on a horizontal or vertical number line diagram.</li> <li>d. Apply properties of operations as strategies to add and subtract rational numbers.</li> <li>7.NS.3. Solve real-world and mathematical problems involving the four operations with rational numbers.</li> </ul>
	M07.A-N.1.1.3 Apply properties of operations to multiply and divide rational numbers, including real-world contexts; demonstrate that the decimal form of a rational number terminates or eventually repeats.	<ul> <li>7.NS.2. Apply and extend previous understandings of multiplication and division and of fractions to multiply and divide rational numbers.</li> <li>c. Apply properties of operations as strategies to multiply and divide rational numbers.</li> <li>d. Convert a rational number to a decimal using long division; know that the decimal form of a rational number terminates in 0s or eventually repeats.</li> <li>7.NS.3. Solve real-world and mathematical problems involving the four operations with rational numbers.</li> </ul>

Updated 5.18.2012 Page 2 of 9



an educational service agency		
Grade 7		
PA Common Core Standard	PA Eligible Content	Common Core State Standard
	NO MATCH	<ul> <li>7.NS.1. Apply and extend previous understandings of addition and subtraction to add and subtract rational numbers; represent addition and subtraction on a horizontal or vertical number line diagram.</li> <li>a. Describe situations in which opposite quantities combine to make 0. For example, a hydrogen atom has 0 charge because its two constituents are oppositely charged.</li> <li>b. Understand p + q as the number located a distance  q  from p, in the positive or negative direction depending on whether q is positive or negative. Show that a number and its opposite have a sum of 0 (are additive inverses). Interpret sums of rational numbers by describing realworld contexts.</li> <li>c. Understand subtraction of rational numbers as adding the additive inverse, p - q = p + (-q). Show that the distance between two rational numbers on the number line is the absolute value of their difference, and apply this principle in real-world contexts.</li> </ul>
	NO MATCH	<ul> <li>7.NS.2. Apply and extend previous understandings of multiplication and division and of fractions to multiply and divide rational numbers.</li> <li>a. Understand that multiplication is extended from fractions to rational numbers by requiring that operations continue to satisfy the properties of operations, particularly the distributive property, leading to products such as (-1)(-1) = 1 and the rules for multiplying signed numbers. Interpret products of rational numbers by describing real-world contexts.</li> <li>b. Understand that integers can be divided, provided that the divisor is not zero, and every quotient of integers (with non-zero divisor) is a rational number. If p and q are integers, then -(p/q) = (-p)/q = p/(-q). Interpret quotients of rational numbers by describing real-world contexts.</li> </ul>

Updated 5.18.2012 Page 3 of 9



Grade 7		
PA Common Core Standard	PA Eligible Content	Common Core State Standard
<b>2.2.7.B.1</b> Apply properties of operations to generate equivalent expressions.	<b>M07.B-E.1.1.1</b> Apply properties of operations to add, subtract, factor, and expand linear expressions with rational coefficients. <i>Example 1: The expression 1/2</i> • $(x + 6)$ <i>is equivalent to 1/2</i> • $x + 3$ . <i>Example 2: The expression 5.3</i> – $y + 4.2$ <i>is equivalent to 9.5</i> – $y$ (or – $y + 9.5$ ). <i>Example 3: The expression <math>4w - 10</math> is equivalent to <math>2(2w - 5)</math>.</i>	<b>7.EE.1</b> Apply properties of operations as strategies to add, subtract, factor, and expand linear expressions with rational coefficients.
2.2.7.B.3  Model and solve real-world and mathematical problems by using and connecting numerical, algebraic, and/or graphical representations.	<b>M07.B-E.2.1.1</b> Apply properties of operations to calculate with numbers in any form; convert between forms as appropriate.  Example: If a woman making \$25 an hour gets a $10\%$ raise, she will make an additional $1/10$ of her salary an hour, or \$2.50, for a new salary of \$27.50 (or $1.1 \times $25 = $27.50$ ).	<b>7.EE.3</b> Solve multi-step real-life and mathematical problems posed with positive and negative rational numbers in any form (whole numbers, fractions, and decimals), using tools strategically. Apply properties of operations to calculate with numbers in any form; convert between forms as appropriate; and assess the reasonableness of answers using mental computation and estimation strategies. For example: If a woman making \$25 an hour gets a 10% raise, she will make an additional 1/10 of her salary an hour, or \$2.50, for a new salary of \$27.50. If you want to place a towel bar 9 3/4 inches long in the center of a door that is 27 1/2 inches wide, you will need to place the bar about 9 inches from each edge; this estimate can be used as a check on the exact computation.
	<b>M07.B-E.2.2.1</b> Solve word problems leading to equations of the form $px + q = r$ and $p(x + q) = r$ , where $p$ , $q$ , and $r$ are specific rational numbers. <i>Example: The perimeter of a rectangle is 54 cm. Its length is 6 cm. What is its width?</i>	<b>7.EE.4</b> Use variables to represent quantities in a real-world or mathematical problem, and construct simple equations and inequalities to solve problems by reasoning about the quantities. <b>a.</b> Solve word problems leading to equations of the form $px + q = r$ and $p(x + q) = r$ , where p, q, and r are specific rational numbers. Solve equations of these forms fluently. Compare an algebraic solution to an arithmetic solution, identifying the sequence of the operations used in each approach. For example, the perimeter of a rectangle is 54 cm. Its length is 6 cm. What is its width?

Updated 5.18.2012 Page 4 of 9



	Grade 7		
PA Common Core Standard	PA Eligible Content	Common Core State Standard	
	M07.B-E.2.2.2 Solve word problems leading to inequalities of the form px + q > r or px + q < r, where p, q, and r are specific rational numbers, and graph the solution set of the inequality.  Example: A salesperson is paid \$50 per week plus \$3 per sale. This week she wants her pay to be at least \$100. Write an inequality for the number of sales the salesperson needs to make, and describe the solutions.  M07.B-E.2.3.1 Determine the reasonableness of an answer(s), or interpret the solution(s) in the context of the problem.  Example: If you want to place a towel bar that is 9 3/4 inches long in the center of a door that is 27 1/2 inches wide, you will need to place the bar about 9 inches from each edge; this estimate can be used as a check on the exact computation.	<ul> <li>7.EE.4. Use variables to represent quantities in a real-world or mathematical problem, and construct simple equations and inequalities to solve problems by reasoning about the quantities.</li> <li>b. Solve word problems leading to inequalities of the form px + q &gt; r or px + q &lt; r, where p, q, and r are specific rational numbers. Graph the solution set of the inequality and interpret it in the context of the problem. For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions.</li> <li>7.EE.3 Solve multi-step real-life and mathematical problems posed with positive and negative rational numbers in any form (whole numbers, fractions, and decimals), using tools strategically. Apply properties of operations to calculate with numbers in any form; convert between form as appropriate; and assess the reasonableness of answers using menta computation and estimation strategies. For example: If a woman making \$25 an hour gets a 10% raise, she will make an additional 1/10 of her salary an hour, or \$2.50, for a new salary of \$27.50. If you want to place a towel bar 9 3/4 inches long in the center of a door that is 27 1/2 inches wide, you will need to place the bar about 9 inches from each edge; this estimate can be used as a check on the exact computation.</li> </ul>	
	NO MATCH	<b>7.EE.2.</b> Understand that rewriting an expression in different forms in a problem context can shed light on the problem and how the quantities in it are related. For example, $a + 0.05a = 1.05a$ means that "increase by 5%" is the same as "multiply by 1.05."	

Updated 5.18.2012 Page 5 of 9



Grade 7		
PA Common Core Standard	PA Eligible Content	Common Core State Standard
1	M07.C-G.2.1.1 Identify and use properties of supplementary, complementary, and adjacent angles in a multistep problem to write and solve simple equations for an unknown angle in a figure.	<b>7.G.5</b> Use facts about supplementary, complementary, vertical, and adjacent angles in a multi-step problem to write and solve simple equations for an unknown angle in a figure.
	M07.C-G.2.1.2 Identify and use properties of angles formed when two parallel lines are cut by a transversal (e.g., angles may include alternate interior, alternate exterior, vertical, corresponding).	NO MATCH
	M07.C-G.2.2.1 Find the area and circumference of a circle. Solve problems involving area and circumference of a circle(s). Formulas will be provided.	<b>7.G.4.</b> Know the formulas for the area and circumference of a circle and use them to solve problems; give an informal derivation of the relationship between the circumference and area of a circle.
	M07.C-G.2.2.2 Solve real-world and mathematical problems involving area, volume, and surface area of two and three-dimensional objects composed of triangles, quadrilaterals, polygons, cubes, and right prisms. Formulas will be provided.	<b>7.G.6.</b> Solve real-world and mathematical problems involving area, volume and surface area of two- and three-dimensional objects composed of triangles, quadrilaterals, polygons, cubes, and right prisms.
figures and describe the	M07.C-G.1.1.1 Solve problems involving scale drawings of geometric figures, including finding length and area.	<b>7.G.1</b> Solve problems involving scale drawings of geometric figures, including computing actual lengths and areas from a scale drawing and reproducing a scale drawing at a different scale.
relationships between them.	M07.C-G.1.1.2 Identify or describe the properties of all types of triangles based on angle and side measure.	NO MATCH
	<b>M07.C-G.1.1.3</b> Use and apply the triangle inequality theorem.	NO MATCH
	M07.C-G.1.1.4 Describe the two-dimensional figures that result from slicing three-dimensional figures. Example: Describe plane sections of right rectangular prisms and right rectangular pyramids.	<b>7.EE.3.</b> Describe the two-dimensional figures that result from slicing three-dimensional figures, as in plane sections of right rectangular prisms and right rectangular pyramids.

Updated 5.18.2012 Page 6 of 9



Grade 7		
PA Common Core Standard	PA Eligible Content	Common Core State Standard
	NO MATCH	<b>7.EE.2.</b> Draw (freehand, with ruler and protractor, and with technology) geometric shapes with given conditions. Focus on constructing triangles from three measures of angles or sides, noticing when the conditions determine a unique triangle, more than one triangle, or no triangle.
2.4.7.B.1 Draw inferences about populations based on random sampling concepts.	M07.D-S.1.1.1 Determine whether a sample is a random sample given a real-world situation.	<b>7.SP.1.</b> Understand that statistics can be used to gain information about a population by examining a sample of the population; generalizations about a population from a sample are valid only if the sample is representative of that population. Understand that random sampling tends to produce representative samples and support valid inferences.
	characteristic of interest.	<b>7.SP.2</b> Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.
2.4.7.B.2  Draw informal comparative inferences about two populations.	Example 1: The mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team. This difference	7.SP.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability (mean absolute deviation) on either team; on a dot plot, the separation between the two distributions of heights is noticeable. 7.SP.4. Use measures of center and measures of variability for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.

Updated 5.18.2012 Page 7 of 9



Grade 7		
PA Common Core Standard	PA Eligible Content	Common Core State Standard
<b>2.4.7.B.3</b> Investigate chance processes and develop, use, and evaluate probability models.	M07.D-S.3.1.1 Predict or determine whether some outcomes are certain, more likely, less likely, equally likely, or impossible (i.e., a probability near 0 indicates an unlikely event, a probability around 1/2 indicates an event that is neither unlikely nor likely, and a probability near 1 indicates a likely event).	<b>7.SP.5</b> Understand that the probability of a chance event is a number between 0 and 1 that expresses the likelihood of the event occurring. Larger numbers indicate greater likelihood. A probability near 0 indicates an unlikely event, a probability around 1/2 indicates an event that is neither unlikely nor likely, and a probability near 1 indicates a likely event.
	event given relative frequency. Predict the	<b>7.SP.6.</b> 6. Approximate the probability of a chance event by collecting data on the chance process that produces it and observing its long-run relative frequency, and predict the approximate relative frequency given the probability. For example, when rolling a number cube 600 times, predict that a 3 or 6 would be rolled roughly 200 times, but probably not exactly 200 times.
	M07.D-S.3.2.2 Find the probability of a simple event, including the probability of a simple event <b>not</b> occurring.  Example: What is the probability of <b>not</b> rolling a 1 on a number cube?	NO MATCH
	M07.D-S.3.2.3 Find probabilities of independent compound events using organized lists, tables, tree diagrams, and simulation.	<ul> <li>7.SP.8. Find probabilities of compound events using organized lists, tables, tree diagrams, and simulation.</li> <li>a. Understand that, just as with simple events, the probability of a compound event is the fraction of outcomes in the sample space for which the compound event occurs.</li> <li>b. Represent sample spaces for compound events using methods such as organized lists, tables and tree diagrams. For an event described in everyday language (e.g., "rolling double sixes"), identify the outcomes in the sample space which compose the event.</li> <li>c. Design and use a simulation to generate frequencies for compound events. For example, use random digits as a simulation tool to approximate the answer to the question: If 40% of donors have type A blood, what is the probability that it will take at least 4 donors to find one with type A blood?</li> </ul>

Updated 5.18.2012 Page 8 of 9



an educational service agency		
Grade 7		
PA Common Core Standard	PA Eligible Content	Common Core State Standard
	NO MATCH	<ul> <li>7.SP.7. Develop a probability model and use it to find probabilities of events. Compare probabilities from a model to observed frequencies; if the agreement is not good, explain possible sources of the discrepancy.</li> <li>a. Develop a uniform probability model by assigning equal probability to all outcomes, and use the model to determine probabilities of events. For example, if a student is selected at random from a class, find the probability that Jane will be selected and the probability that a girl will be selected.</li> <li>b. Develop a probability model (which may not be uniform) by observing frequencies in data generated from a chance process. For example, find the approximate probability that a spinning penny will land heads up or that a tossed paper cup will land open-end down. Do the outcomes for the spinning penny appear to be equally likely based on the observed frequencies?</li> </ul>

Updated 5.18.2012 Page 9 of 9